

## **New distributional data of the Turkish Ichneumonidae (Hymenoptera) II. Ctenopelmatinae and Campopleginae**

Hikmet ÖZBEK\*      Saliha PEKEL\*      Janko KOLAROV\*\*

\* Atatürk University, Agriculture Faculty, Department of Plant Protection, 25240  
Erzurum, TURKEY, e-mail: hozbek@atauni.edu.tr

\*\*University of Sofia, BULGARIA

### **ABSTRACT**

Faunistical and distributional data for 17 species of the subfamilies Ctenopelmatinae and Campopleginae from Turkey are given. Of these, *Absyrtus vernalis* Bauer, *Bathylectes anurus* Thomson, *Charops cantator* De Geer, *Dusona ucrainica* Hinz and *D. contumax* Förster are new records for the Turkish fauna. However, *Alcima pictor* Aubert is endemic in Turkey.

*Key words:* Ichneumonidae, Ctenopelmatinae, Campopleginae, Turkey, fauna.

### **INTRODUCTION**

After the subfamilies Pimplinae and Tryphoninae treated in the first part of our serial studies (Kolarov et al., 1999), the present paper deals with the subfamilies Ctenopelmatinae and Campopleginae of the family Ichneumonidae. The vast majority of the ctenopelmatines are koinobiont endoparasitoids of larval tentredinoid and megalodontoid sawflies (Gauld & Bolton, 1988). Very few species parasitize lepidopterous larvae (Heath, 1961). Campoplegines are solitary koinobiont endoparasitoids and the majority attack Lepidoptera, a smaller number of species parasitize tentredinoid sawflies and some phytophagous Coleoptera (Gauld & Bolton, 1988).