

Two new names for the genus *Amarygmus* and one replacement for the genus *Strongylium* (Coleoptera: Tenebrionidae)

Vratislav Richard E.M.J.-B. Bejsak-Collorodo-Mansfeld

The Wentworth Bldg., B.62 : University of Sydney 2006 : New South Wales

e-mail: vratislav@bigfoot.com

Carter (1913) described new Tenebrionidae from Australia which included two new species of *Amarygmus*, *A. carbo* from Kuranda and the Bloomfield River and *A. femoratus* from Port Darwin.

Kaszab (1956, 1980) was apparently unaware that these names had been used in *Amarygmus*. Kaszab (1956) described *Amarygmus femoratus* from the Cyclops Mountains, New Guinea, and later (Kaszab 1980) described *A. carbo* from Ceylon. I propose new names for Kaszab's species I propose new names for Kaszab's species: *Amarygmus femoratus* Kaszab, 1956 = *A. lawrenci* (nomen novum) and *Amarygmus carbo* Kaszab, 1980 = *A. masumoto* (nomen novum).

Recently I (B.-C.-M. 1999) accidentally created another synonym, *Strongylium becvari*, when proposing a replacement name for *S. anthrax* Kaszab, (1977) the name *S. becvari* having already been used by Masumoto (1998). I propose *Strongylium stanislavi* Bejsak-Collorodo-Mansfeld, nomen novum, to replace both Kaszab's and my synonyms.

REFERENCES

- Carter, H.J. 1913. Notes and tabulation of the Australian *Amarygmus* (family Tenebrionidae), with descriptions of new species. *Transactions and Proceedings of the Royal Society of South Australia* 37: 6-47.