

***Spiximorpha worelli* (Brădescu, 1972), (Diptera: Syrphidae) New
Record for the Fauna of Turkey**

Süleyman SARIBIYIK

Gazi University, Kastamonu Education Faculty, 37200 Kastamonu, TURKEY

e-mail: sbiyik@gazi.edu.tr

ABSTRACT

Spiximorpha worelli (Brădescu, 1972) is the first record for the Turkish *Syrphidae* fauna. The specimen was collected from Kastamonu province.

Key words: *Spiximorpha worelli*, new record, Turkey

INTRODUCTION

Only one species of *Spiximorpha* Rondani, 1850, (*Cerioides*, Rondani, 1850), (*Cerioides euprosopa*, Loew, 1869) has been reported from Turkey so far. This species was recorded for the first time from Turkey (İzmir) by Sack (1932) and Peck (1988).

Spiximorpha worelli (Brădescu, 1972), reported in the present paper, has been identified by the author. Sack (1932) and (Brădescu, 1972) were used for identification.

In this work, the faunistic reports of the species are presented along with diagrams of body parts showing important distinguishing features.

Spiximorpha worelli (Brădescu, 1972) is deposited in the Zoological Museum of Gazi University (ZMGU), Arts & Sciences Faculty, Ankara, Turkey.