

**Additional Notes to the Knowledge of Longhorned Beetle
Collection from Zoological Museum of Gazi University, Ankara,
Turkey (ZMGU) for Turkish Fauna (Coleoptera, Cerambycidae)**

Hüseyin ÖZDIKMEN

Ersin DEMIREL

Gazi University, Science and Arts Faculty, Department of Biology, 06500 Ankara,
TURKEY, e-mail: ozdikmen@gazi.edu.tr

ABSTRACT

In the present paper specimens of the family Cerambycidae collection (Coleoptera) in ZMGU, Ankara/Turkey. New data are presented on Cerambycidae fauna of Turkey were studied. The faunistic data in the present paper on almost all species add to the knowledge of their distribution in Turkey. As a result of identification of these specimens sixty four species and two subspecies belonging to thirty six genera of four subfamilies for Turkey were determined. As mainly interesting records: *Anoplodera sexguttata* (Fabricius, 1775) is rare taxon for Turkey. *Pedostrangalia verticenigra* (Pic, 1892), *Clytus ciliciensis* Chevrolat, 1863 and *Clytus taurusiensis* (Pic, 1913) are fairly local and endemic taxa for Turkey. Also, *Clytus schurmanni* Sama, 1996 and *Oberea erythrocephala schurmanni* Heyrovsky, 1962 are known to occur only in Turkey. Furthermore, many taxa are new for various provinces of Turkey.

Key words: Prioninae, Lepturinae, Cerambycinae, Lamiinae, Cerambycidae, Coleoptera, Turkey, ZMGU

INTRODUCTION

Especially since last century, the studies which were mainly as faunistic and taxonomic works, on longhorn beetles fauna of Turkey have been carried out. The chief faunistic studies are Fairmaire (1884); Semenov (1900); Bodemeyer (1900, 1906); Ýriboz (1938, 1940); Acatay (1943, 1963); Schimitschek (1944); Alkan (1956); Erdem (1947, 1968); Defne (1964); Çanakçýođlu (1956, 1993); Villiers (1959, 1967); Demelt & Alkan (1962); Demelt (1963, 1967); Nizamlyođlu & Gökmen (1964); Perissinotto & Luchini (1966); Breuning & Villiers (1967); Besçeli (1969); Fuchs & Breuning (1971); Ekici (1971); Gfeller (1972); Tuatay *et al.*,